

INCIDENCE AND ECOLOGY OF MARINE FOULING ORGANISMS  
IN THE EASTERN HARBOUR OF ALEXANDRIA, EGYPT.

M. MOHAMED EL-KOMI\*

\* National Institute of Oceanography and Fisheries,  
Alexandria.

ABSTRACT

The seasonal changes, intensity and constituents of fouling developed on exposed test panels for short and long term intervals are studied in relation to the prevailing environmental conditions in the Eastern Harbour of Alexandria, from March 1983 to March 1984. The intensity of fouling on exposed test panels for long durations is considerably more dense than the total fouling developed on exposed panels for short term successively during the same period. The fouling colonized on submerged panels for long intervals reaches to a "Saturation Point" after 3 to 6 successive months.

INTRODUCTION

The Eastern Harbour of Alexandria is relatively a small semi-circular polluted bay, covering an area of about 2.8 km<sup>2</sup>. It receives many kinds of vessels specially fishing boats. In addition, a large amount of untreated sewage flows into it amount about 36.000 m<sup>3</sup>/day (Said and Maiyza 1987). The environmental conditions, in this area are greatly variable. The fouling communities in the Eastern Harbour of Alexandria were reported by Banoub (1960), Megally (1970) and Ghobashy (1976). The present work was conducted to demonstrate the seasonal variations, intensity and constituents of attachment of fouling organisms on exposed test panels for short and long durations of time. The occurrence of main stages of foulers in sea water was indicated in relation to the prevailing environmental parameters in the Harbour to give an idea about the respective periods of immersion. It is possible to make interesting successional growth and longevity of fouling groups inhabited the submerged objects.

MATERIALS AND METHODS

Observations were made during the period from March 1983 to March 1984 at a raft located in the Eastern Harbour of Alexandria. The fouling organisms were collected monthly by exposing test panels made of polystyrene (15 x 15 cm) fixed to an iron frame sized 1 x 0.8 m in two rows. This frame was

immersed vertically from the raft at about 1.5 m below the sea surface. The panels were replaced and taken regularly at the beginning of every month. The exposed test panels were arranged in two series, a short term for one month exposure period and a long term for various intervals of immersions from two months to one year successively. The settling number of organisms, wet weight of the principal fouling groups and the growth of organisms with the knowledge of the approximate time of settlement on test panels were carried out. Water samples were taken monthly from the surface and 4.5 m depth. These samples were used for the measurements of salinity, dissolved oxygen, oxidizable oxygen and alkalinity according to Strickland and Parsons (1968). Plankton hauls were taken to examine the occurrence of the pelagic stages of common fouling organisms in the study area.

## RESULTS

### 1- Environmental Conditions:

The physical and chemical data are illustrated in Fig. 1. The annual fluctuation in temperature of sea water ranged from 28°C in August 1983 (summer) to 16°C in February 1984 (winter). On the other hand, the fluctuation in the values of pH, dissolved oxygen and salinity of sea water throughout most of the year was not too great.

### 2- Seasonal Changes in the Fouling Populations: Short-term exposure panels

The fouling developed on the submerged panels for one month duration was very little both in settlement density and growth. A total of 34 different kinds of fouling were found to settle every month of the year. These are listed in Table 1. The general fouling picture will be presented seasonally rather than monthly.

#### a- Spring fouling (March-May). The average water temperature was 22°C.

During this period the fouling developed on panels every month was relatively poor. Algae were predominated by *Ulva intestinalis*, *Ectocarpus conferoides*, *Cladophora* sp., and *Enteromorpha* sp. The growth of these plants reached 11, 4, 8 and 13.4 cm long after one month of settlement, respectively. While, *Bugula neritina* encountered in large numbers during April and May 1983 with an average of 118 col/100 cm<sup>2</sup>, attained small size from 2 to 9 bifurcations.

The plankton samples included mostly of *Polychaeta*, barnacles and *Bugula* larvae (Table 2) during May 1983 were encountered by 1026, 1944 and 108 larvae/m<sup>3</sup>, respectively.

#### b- Summer fouling (June-August). The average water temperature was 27°C.




FIG. 1  
 Seasonal variations of environmental factors in the  
 Eastern Harbour of Alexandria at the surface (----)  
 and at depth of 4.5 m below the sea surface (—),  
 during March 1983 to March 1984.

Table 1

Monthly changes in the settlement of fouling organisms on panels (No. org/100 cm<sup>2</sup>) and the total biomass during March 1983 to February 1984 in the Eastern Harbour of Alexandria

Species	May 1983	Apr	May	Jun	Jul	Aug	Sep	Oct	Nov	Dec	Jan 1984	Feb
<b>ALGAE</b>												
<i>Ulva intestinalis</i>	-	C	C	P	-	-	P	P	P	A	A	P
<i>Cladophora</i> sp.	P	-	-	-	-	-	-	-	-	P	P	C
<i>Chaetomorpha</i> sp.	-	-	-	-	-	C	C	-	-	-	-	-
<i>Enteromorpha</i> sp.	A	-	-	P	-	-	-	-	-	-	C	P
<i>Ectocarpus</i> sp.	C	C	C	C	C	C	C	A	A	A	C	C
<i>Ceramium</i> sp.	-	-	-	-	-	-	-	P	P	P	-	P
<b>HYDROIDS</b>												
<i>Obelia geniculata</i>	-	-	C	-	P	C	C	C	C	P	C	P
<i>Tubularia larynx</i>	-	5	-	-	-	-	-	-	-	7	11	-
<b>BRYOZOA</b>												
<i>Bugula neritina</i>	10	118	88	41	11	-	-	4	28	26	4	66
<i>Bugula turbinata</i>	-	-	-	-	-	-	-	-	-	-	-	-
<i>Zoobotryon</i> sp.	-	-	-	-	-	-	C	P	P	-	-	-
<i>Boerbankia</i> sp.	-	-	-	-	-	-	-	-	-	P	-	-
<i>watersipora</i> sp.	-	-	-	-	-	-	-	-	-	P	-	-
<b>POLYCHAET</b>												
<i>Hydriodes elegans</i>	-	5	396	5280	352	83	379	4400	2640	1012	154	29
<i>Serpula vermicularis</i>	-	88	P	P	P	P	P	P	P	P	P	P
<i>Spirorbis</i> sp.	-	-	-	-	-	4	-	-	-	-	-	-
<i>Syllis</i> sp.	-	-	-	-	4	5	-	9	6	8	-	-
<i>Nereis diversicolor</i>	-	1	18	4	5	7	-	14	4	-	-	3

Table 1 (Continued)

	308	792	704	1100	103	154	88	253	1100	1012	88	88
<b>BARINACLES</b>												
Balanus amphitrite	1	1	1	7	4	4	5	12	36	10	11	16
Balanus eburneus	-	-	-	2	1	1	-	1	-	2	-	-
Balanus perforatus	-	4	2	32	2	2	-	-	-	-	1	9
Balanus trigonus	-	-	-	1	1	-	-	-	-	-	-	-
Amphipod tubes	308	792	704	1100	103	154	88	253	1100	1012	88	88
Corophium sextomi	4	46	28	23	6	8	37	70	79	100	15	15
Erichthoneus brasii	42	238	370	286	-	7	5	8	-	37	-	8
Jassa falcata	210	8	17	66	3	-	-	-	-	-	-	3
Stenothoe sp.	26	-	-	21	11	14	-	33	11	27	8	4
Elasmopus pectenarius	-	-	-	117	-	-	17	26	59	52	40	9
Caprella equilibra	-	6	6	2	-	-	-	-	-	-	-	-
Tanaia cavolinii	2	5	4	2	1	-	7	23	40	18	10	5
Idotea baltica	2	4	1	-	-	-	-	-	-	-	-	-
Cymodoce truncata	-	-	1	-	-	-	-	-	-	-	-	-
Dynamene bidentata	-	-	-	37	-	-	-	-	-	-	-	-
<b>ASCIDIANS</b>												
Diplosoma listerianum	-	-	1	-	-	-	-	-	-	-	-	-
Botryllus schlosseri	-	10	-	-	-	-	-	-	-	-	-	-
<b>Platyhelminthes</b>												
	-	-	-	10	-	-	-	-	-	-	-	-
Total wet weight	4	8	6	38	4	3	2	6	7	13	2	5

A = Abundant, C = Common, and P = Present

Table 2

Seasonal changes of the planktoic larvae in the Eastern Harbour Alexandria (No. of larvae /m<sup>3</sup>) recorded in the vertical hauls during the period from March 1983 to March 1984.

Groups	Mar 1983	Apr	May	Jun	Jul	Aug	Sep	Oct	Nov	Dec	Jan 1984	Feb	Mar
Leptomed- usae	-	-	-	114	57	-	22	69	360	-	220	19	120
Bugula spp. larvae	-	-	108	-	-	-	-	-	-	-	66	-	-
Polychaeta larvae	420	600	1026	7410	2679	1508	1496	1172	168	825	290	230	2112
Barnacles Nauplius l.	240	300	1944	798	342	884	220	414	264	2585	110	38	1248
Barnacles cypria l.	20	120	-	57	-	156	-	-	-	-	-	-	-
Bivalve larvae	-	60	-	114	-	-	1385	161	-	-	44	-	-
Gastropod larvae	-	-	-	-	-	-	-	-	-	185	22	-	-
Ascidians larvae	20	20	-	-	-	-	-	-	-	-	22	-	48
Appendicu- larie	-	-	-	228	798	156	-	115	-	55	-	-	-
Total No. org/m <sup>3</sup>	700	1100	3078	8721	3876	2704	3123	1931	792	3630	774	287	3528

The fouling was also very poor. It was relatively more dense on the June panel weighed an average of 38 gm/100 cm<sup>2</sup>. It embraced mainly calcareous tube worms weighed 27 gm. Alga, *Ectocarpus conferoides* dominated during summer months whereas *Chaetomorpha* sp. was common on August panel. The growth of algae, on the other hand, was very little about one cm long.

The majority of plankton samples of Polychaet, Barnacles larvae and Leptomedusae, reached more than 7410, 855 and 114 org./m<sup>3</sup>, respectively.

C- Autumn fouling (September-November).  
The average water temperature was 23°C.

The quantity and quality of fouling colonized on panels during Autumn months were also very poor. The wet weight of fouling was about 7 gm/100 cm<sup>2</sup>. It consisted mainly of algae (*Ectocarpus conferoides* and *Ulva intestinalis*), tube worms (*Hydriodes elegans* and *Serpula vermicularis*) and Hydriods

(*Obelia geniculata*) while, *Balanus amphitrite* is the only barnacle species settled during this season and encountered in few number (5-36 ind./100 cm<sup>2</sup>) attained sexual maturity at 8 mm in basal diameter. Polychaet, Barnacles and leptomedusae were still abundant in the plankton samples numbered 1500, 1300, and 530 org./ m<sup>3</sup>, respectively. Bivalve veliger larvae dominated during September (1385 org./m<sup>3</sup>).

d- Winter fouling (December 1983-February 1984).  
The water temperature decreased to 16°C in January.

The fouling assemblaged during winter months was represented by about 3-13 gm/100 cm<sup>2</sup>, and contained large quantity of algae, *Ulva intestinalis* and *Ectocarpus conferridoides* rather than *Cladophora* sp. and *Ceramium* sp. The other fouling included few colonies of Hydriods (*Obelia geniculata* and *Tubularia larynx*), Bryozoa (*Bugula neritina* and *B. turbinata*), and few individuals of tube worms and barnacles were recorded.

Barnacles larvae yielded high values during December (2585 org./m<sup>3</sup>). It does not reflect the real attachment numbers of barnacles grown on the month panel. This may be due to low temperature during December.

### 3- Species composition and settlement of fouling organisms on Long-term exposure panels

During the investigation period from March 1983 to March 1984, a total of 41 species of sedentary marine fouling organisms were recorded on the long term panels (Table 3). These species belonged to 8 main fouling groups namely, Barnacles, Serpulid, Ascidians, Bryozoa, Amphipod mud building tubes, Hydriods, Algae and Sponges. They are considered among the main constituents of fouling organisms growing in many harbours of the world, in addition to 21 species of free-living forms; four Polychaets, five Decapods, seven Amphipods, five Isopods and unidentified Platyhelminthes species.

a- Barnacles were the predominant population and the most persistent fouling grown on the submerged surfaces especially for prolonged durations of immersion (Fig. 2 & Table 4). They showed maximum growth rate during the first two months (Table 5) and their survival extended for 8 successive months or more. They were represented by four species namely, *Balanus amphitrite*, *B. eburneus*, *B. perforatus* and *B. trigonus*.

b- Serpulid calcareous tube worms were numerically the most abundant animals which were attached in considerable amount forming a dense mat, usually reaching a thickness of more or less 3 cm. They flourished well on the panels exposed during spring and early summer months for longer intervals (Fig. 2 and Tables 4 & 5). This may be attributed to the breeding seasons and the overcrowded tube worms can be easily removed under the effect of the external circumstances. This

Table 3

Species composition of fouling recorded on long-term panels during various durations of immersion from March 1983 to February 1984, in the Eastern Harbour of Alexandria.

---

<b>Algae</b>	
Ulva intestinalis	Enteromorpha sp.
Chaetomorpha sp.	Cladomorpha sp.
Polysiphonia sp.	Codium sp.
Ectocarpus sp.	Ceramium sp.
<b>Sponges</b>	
<b>Hydriods</b>	
Obelia geniculata	Lubularia larynx
<b>Bryozoa</b>	
Bugula neritina	Bugula turbinata
Zoobotryon sp.	Bowerbankia gracilis
Bowerbankia imbricata	Cryptosula pallasiana
Schizoporella errata	
<b>Platyhelminthis</b>	
<b>Polychaeta</b>	
Hydriodes elegans	Hydriodes dianthus
Hydriodes dirampha	Serpula vermicularis
Pomatoceros triqueter	Dasychone sp.
Polydora sp.	Eupolyornia sp.
Nereis diversicolor	Scale worms
<b>Barnacles</b>	
Balanus amphitrite	Balanus eburneus
Balanus perforatus	Balanus trigonus



Table 3 (Continued)

---

Amphipod building tubes	
Amphipod	
Corophium sextomi	Erichthonius brasiliensis
Jassa falcata	Stenothoe sp.
Elasmopus pectenía	Caprella equilibra
Tanais cavolinii	
Isopod	
Cirolana aegyptica	Cymodoce truncata
Dynamene bidentus	Sphaeroma sp.
Idotea baltica	
Decapod	
Pachygrapsus sp.	Xantho sp.
Eriphia sp.	Thia sp.
Alepheus sp.	
Mollusca	
Venerupus sp.	Cardium sp.
Anomia ephippium	
Ascidians	
Ciona intestinalis	Ascidia mentula
Styela partita	Styela plicata
Diplosoma listerianum	Botryllus schlosseri
Botrylloides sp.	

---




FIG. 2  
 Seasonal variations of fouling biomass (wet weight/100 cm<sup>2</sup>) developed on submerged test panels monthly and bimonthly (the upper half) and long term exposure periods from 3 to 12-months (the lower half) during March 1983 to February 1984 in the Eastern Harbour of Alexandria. Numbers illustrate the immersion durations in months.

Table 4

Wet weight of main fouling groups developed on panels in grams per 100 cm<sup>2</sup> for variable immersion intervals during March 1983 to February 1984 in the Eastern Harbour of Alexandria.

Duration months	Period of immersion	Algae	Hydrifods	Bryozoa	Serpulids	Barnacles	Ascidians	Other groups	Total wet weight
2	Mar-Apr	3	+	4	30	1	23	2	63
2	Jul-Aug	+	+	+	13	7	-	2	22
2	Sep-Oct	+	+	+	4	1	-	1	6
2	Oct-Nov	1	+	2	1	2	1	1	8
2	Nov-Dec	2	1	3	5	1	-	1	13
2	Dec-Jan	+	-	1	9	4	-	1	15
2	Jan-Feb	2	-	1	2	2	-	1	8
3	Mar-May	-	1	27	56	1	93	1	179
3	Jun-Aug	-	-	+	54	20	6	1	81
3	Sep-Nov	2	-	3	1	5	-	1	12
3	Oct-Dec	-	+	7	12	17	1	1	38
3	Nov-Jan	+	+	1	18	13	1	1	34
3	Dec-Feb	+	-	6	29	17	2	1	55
4	Mar-Jun	-	-	78	119	4	5	3	209
4	Jun-Sep	-	+	+	29	83	14	2	128
4	Sep-Dec	+	4	+	3	10	1	1	19
4	Oct-Jan	-	-	47	29	41	4	1	122
5	Mar-Jul	+	-	5	99	17	3	4	128
5	Apr-Aug	-	+	-	6	31	3	1	41
5	Oct-Feb	+	+	13	4	39	4	1	61
6	Mar-Aug	-	+	+	28	20	3	1	52
6	Jun-Nov	-	-	3	7	145	6	2	163
7	Mar-Sep	-	+	+	169	46	11	3	229
8	Mar-Oct	+	4	+	8	94	5	1	112
9	Mar-Nov	+	+	9	5	125	19	2	160
10	Mar-Dec	+	+	33	7	101	32	2	175
11	Mar-Jan	+	+	22	10	74	64	1	171
12	Mar-Feb	-	-	53	20	98	71	1	243

+ Present in a few numbers; - Not observed

Table 4 (Continued)

Duration of Months	Period of immersion	Serpulids *		Ascidians **			
		Hydriodes elegans	Pomatoceros triqueter	Ciona intest- inalis	Styela partita	Styela plicata	Ascidia mentula
2	Mar-Apr	2775(3.5)	-	134(4.2)	-	-	-
2	Jul-Aug	770(2.8)	-	-	-	-	-
2	Sep-Oct	685(1.2)	-	-	-	-	-
2	Oct-Nov	730(1.8)	3(1.1)	-	4(0.6)	-	-
2	Nov-Dec	760(1.5)	-	-	-	-	-
2	Dec-Jan	1375(1.8)	-	-	-	-	-
2	Jan-Feb	190(1.4)	-	-	-	-	-
3	Mar-May	3990(2.7)	-	19(9.5)	1(0.5)	-	-
3	Jun-Aug	3550(3.2)	-	1(4.8)	12(1.6)	1(3.5)	-
3	Sep-Nov	880(2.2)	5(1.6)	-	11(0.4)	-	-
3	Oct-Dec	1440(2.5)	-	-	8(0.8)	-	-
3	Nov-Jan	1665(2.0)	-	1(4.0)	1(4.0)	-	-
3	Dec-Feb	2660(2.0)	-	1(4.0)	-	-	1(3.2)
4	Mar-Jun	8430(3.5)	-	-	-	-	-
4	Jun-Sep	2660(2.5)	3(2.0)	1(4.0)	18(1.4)	3(1.0)	1(3.2)
4	Sep-Dec	550(1.4)	1(2.0)	1(2.5)	9(1.0)	-	-
4	Oct-Jan	2440(2.0)	-	1(4.5)	12(0.8)	1(0.6)	1(1.4)
5	Mar-Jul	7100(3.0)	-	1(2.8)	6(1.2)	-	-
5	Apr-Aug	710(2.6)	1(2.8)	1(7.0)	6(1.2)	-	-
5	Oct-Feb	710(1.8)	-	1(2.0)	-	-	-
6	Mar-Aug	2230(3.0)	-	1(5.0)	5(1.4)	1(3.5)	-
6	Jun-Nov	1330(1.8)	4(1.2)	-	14(1.0)	1(1.2)	1(3.0)
7	Mar-Sep	15760(3.0)	9(2.0)	2(5.0)	14(1.0)	2(1.2)	1(1.2)
8	Mar-Oct	1120(1.2)	-	1(7.5)	5(0.8)	1(0.8)	-
9	Mar-Nov	770(2.2)	12(1.4)	1(2.2)	28(1.0)	3(1.4)	1(3.5)
10	Mar-Dec	790(2.2)	4(1.6)	1(4.5)	54(1.0)	3(2.6)	3(1.8)
11	Mar-Jan	710(2.2)	-	1(6.5)	58(1.4)	4(0.5)	5(0.3)
12	Mar-Feb	3100(2.5)	-	1(4.0)	88(1.8)	4(3.0)	3(3.2)

\* Serpulids: length of tube worms; \*\* Ascidians: length of the individual.

Table 5

Numbers of common foulers developed on panels per 100 cm<sup>2</sup> and their maximum size (cm) appears between brackets, for variable immersion intervals during March 1983 to February 1984 in the Eastern Harbour of Alexandria.

Duration of Months	Period of immersion	Bryozoa *			Barnacles **		
		Bugula neritina	Bugula turbinata	Balanus amphitrite	Balanus eburneus	Balanus perforatus	Balanus trigonus
2	Mar-Apr	24(11)	5(7)	3(0.5)	-	5(0.6)	-
2	Jul-Aug	1(2)	-	5(1.1)#	1(1.3)#	1(1.0)#	1(0.8)
2	Sep-Oct	1(9)#	-	3(1.0)	2(1.4)#	-	1(0.5)
2	Oct-Nov	1(8)	-	5(1.2)#	4(1.2)#	1(0.6)	1(0.4)
2	Nov-Dec	6(10)#	1(4)	5(0.7)	1(0.4)	2(0.5)	1(0.7)
2	Dec-Jan	19(8)	3(8)	17(0.6)	5(0.5)	1(0.4)	-
2	Jan-Feb	21(9)	4(10)	21(0.6)	1(0.5)	27(0.5)	-
3	Mar-May	13(20)#	2(7)	1(0.5)	-	32(0.7)	-
3	Jun-Aug	3(6)	1(4)	25(1.4)#	8(1.2)#	16(1.7)#	6(1.0)#
3	Sep-Nov	4(14)#	1(5)	17(1.2)#	9(1.2)#	2(0.5)	1(0.6)
3	Oct-Dec	8(16)	1(6)	3(1.3)#	13(1.4)#	6(1.1)	6(1.0)
3	Nov-Jan	8(8)	4(9)	34(1.2)#	12(1.3)#	15(1.2)	11(1.4)
3	Dec-Feb	14(12)	24(16)	43(1.0)	30(1.2)	19(0.7)	-
4	Mar-Jun	50(20)#	1(4)	4(0.6)	1(0.6)	24(1.1)#	2(0.9)
4	Jun-Sep	6(4)	1(3)	57(1.2)#	36(1.4)#	17(1.1)#	12(1.0)#
4	Sep-Dec	4(5)	1(4)	16(1.2)#	8(1.4)#	2(1.0)#	4(1.2)#
4	Oct-Jan	40(>20)#	14(15)#	32(1.2)#	14(1.4)#	10(1.2)#	8(1.2)
5	Mar-Jul	5(15)#	2(9)	5(0.8)#	1(1.0)#	79(1.0)#	-
5	Apr-Aug	-	-	41(1.2)#	13(1.2)#	15(1.3)#	9(1.3)#
5	Oct-Feb	10(<20)#	7(18)#	37(1.2)	16(1.2)	14(1.2)	10(1.7)
6	Mar-Aug	2(6)	-	40(1.0)#	12(1.6)	18(1.3)#	5(1.0)#
6	Jun-Nov	3(16)#	3(9)#	36(9)#	42(1.6)	50(1.3)#	18(0.8)#
7	Mar-Sep	5(4)	1(3)	32(1.0)#	15(1.4)#	38(1.2)#	6(1.3)#
8	Mar-Oct	3(14)#	1(4)	14(1.3)#	18(1.4)#	24(1.6)#	22(1.0)#
9	Mar-Nov	4(15)#	3(10)#	104(1.3)#	48(1.8)#	25(1.7)#	12(1.2)#
10	Mar-Dec	21(20)#	4(12)#	93(1.3)	30(1.6)	21(1.4)	19(1.2)
11	Mar-Jan	14(20)#	10(18)#	47(1.6)	32(1.8)	21(1.5)	17(1.3)
12	Mar-Feb	36(<20)#	46(20)#	91(1.6)	54(1.8)	47(2.0)	24(1.6)

\* Bryozoa: Np. of bifurcations; \*\* Barnacles: Rostro-carinal diameter (cm)  
# Individuals reached sexual maturity.

community included 6 species of Serpulids (Table 3), predominated by *Hydroids elegans* and *Serpula vermicularis*. The other four species were found in few numbers.

c- The solitary Ascidians *Ciona intestinalis*, *Styela partita*, *S. plicata* and *Ascidia mentula* grew well on panels immersed for long periods. The first species dominated the bimonthly panels. *S. partita* and *S. plicata* could survive for 6 to 8 successive months (Table 5).

d- *Bugula neritina* and *B. turbinata* were the main erect Bryozoa prevailed on the submerged test panels for long term. They appeared in large numbers of colonies reaching more than 20-bifurcations (Table 5) and could survive for about 4 successive months. They developed at the same time of the development of algae on the exposed panels except during summer months. Two species of encrusting and stolonate bryozoans species were recorded on the long term panels (Table 3).

e- Amphipod building tubes were found in large quantities during the year. The densely fouled panels were more favourable for the formation of these muddy tubes in particular on long exposure panels during summer and autumn seasons. The predominance of amphipod species is as follows: *Corophium sextoni* and *Elasmopus pectenis* outnumbered both *Erichthonius barasilie* and *Stenothoe* sp. while each of *Jassa falcata*, *Tanias cavolinii* and the tubuleless *Caprella equilibra* were less frequent.

f- Hydroids occurred in few quantities on test panels during most of the year, especially on prolonged exposure panels. Colonies of *Obelia geniculata* were outnumbered by *Tubularia larynx*. The latter species was found during April, May and December 1983 after 2, 3 and 10 months immersion, respectively.

g- Sponges were found in very little numbers on long term panels. They occurred on November, 1983 panels after 3, 6 and 9 months and on January, 1984 panel after 3 months immersion. The colony attained size about 8 to 22 mm long.

h- Free-living fouling organisms were represented by small numbers in particular on prolonged test panels almostly during the year. They included 3 species of Polychaet, 5 species of Isopod (Table 3).

i- Platyhelminthes were found on panels immersed for long periods and counted in few number inhabited the shells of barnacles especially on June, 1983 panels after one and 4 months immersion.

j- Algae appeared on the submerged panels during most of the year, except in summer months. Flora was mainly belonging to green algae.

## DISCUSSION

pollution caused by draining 36.000 m<sup>3</sup>/day of untreated sewage into the Harbour obviously results seasonal fluctuations in the environmental conditions in particular the dissolved oxygen, oxidizable oxygen and alkalinity of seawater. The presence of fouling larval stages in plankton samples through most of the year did not demonstrate the real settlement intensity of fouling grown on monthly exposed panels especially during the hot months. Therefore, the larvae were prevented from reaching the attachment stage or further growth smothered.

The fouling complex on the exposed test panels tends to be dominated by barnacles, tube worms, bryozoans and ascidians on which barnacles community firstly settled and was sequented by tube worms, colonial and solitary ascidians and erect Bryozoa. The fouling colonized on panels submerged for long durations was considerably more dense than the total fouling collected on short term panels exposed successively during the same period. According to Meadows (1969) the correlation between short and long term fouling biomass is 1: 2: 3 after one, two and three months exposure periods, respectively. This condition may apply under the same conditions of growth, life span and the density of fouling. The fouling constitution on exposed panels for long term reach a "saturation point" after 3 to 6 successive months and some fouling communities attained a dominant position. The same was recorded in Lake Timsah, Suez Canal where the panels reached a saturated point after 6 or 9 months depending on the time of immersion (El-Komi, 1980 and Ghobashy & El-Koim 1980). New attachment of barnacles larvae have to be prevented to make highly fouled surfaces with adult barnacles (Connell, 1961). The growth of fouling organisms was reduced during the winter where the animals grow rapidly during warm seasons even the fouling density is high. Therefore, the water temperature detected the distribution of marine animals and changed abundance in different seasons (Connell, 1974). In this study, however, large seasonal changes in water conditions directly affected the settlement and growth of fouling organisms. On the other hand, the occurrence of large amount of larvae in seawater and the variety of fouling organisms grown on exposed panels for one month period suggest that adult animals fail to reach maturity during short period.

## REFERENCES

- Banoub, M., 1960. Notes on the fouling of glass plates submerged in the Eastern Harbour of Alexandria. *Alex. Inst. Hydrobiol.*, 64: 1-17.
- Connell, J.H., 1961. Effects of competition predation by *Thais lapillus* and other factors on natural populations of barnacles *Balanus balanoides*. *Ecol. Mon.* 31: 61-104.
- Connell, J.H., 1974. Ecology: Field experimental in marine ecology. In: *Experimental Marine Biology*. (ed) Richard N. Mariscal, Academic press, New York and London, 373 p.

- El-Komi, M.M., 1980. Ecological studies on the marine fouling of the Southern Region of the Suez Canal. M. Sc. Thesis University of Mansoura Egypt 151 p.
- Ghobashy, A.F.A., 1976. Seasonal variation and settlement of behaviour of the principal fouling organisms in the Eastern Harbour of Alexandria. Proc. 4 th Intern. Conger. Mari. Corr. Fouling: 213-220.
- Ghobashy, A.F.A, and M. M. El-Komi, 1980. Fouling in the Lake Tamsah (Egypt). Hydrobiol. Bull. Amsterdam, 14 (3): 179-185.
- Meadows, P.S., 1969. Sublittoral fouling communities on northern coast of Britian. Hydrobiologia, 34: 273-294.
- Megally, A.H., 1970. Ecological study on marine fouling organisms in the Eastern Harbour of Alexandria. M. Sc. Thesis, University of Alexandria, 250 p.
- Said A.M. and I. A. Maiyza, 1987. Effect of the domestic sewage discharge on the hydrographic regime of the Eastern Harbour of Alexandria. Bull. Inst. Oceanogr. & Fish. ARE, 13 (2): 1-20.
- Strickland, J.D.H. and T. R. Parsons, 1968. A practical handbook of sea water analysis. 311 p.